arXiv:1203.1381v2 [math.NA] 5 May 2012

CONVERGENCE OF GOAL-ORIENTED ADAPTIVE FINITE ELEMENT METHODS FOR SEMILINEAR PROBLEMS

MICHAEL HOLST, SARA POLLOCK, AND YUNRONG ZHU

ABSTRACT. In this article we develop convergence theory for a class of goal-oriented adaptive finite element algorithms for second order semilinear elliptic equations. We first introduce several approximate dual problems, and briefly discuss the target problem class. We then review some standard facts concerning conforming finite element discretization and error-estimate-driven adaptive finite element methods (AFEM). We include a brief summary of *a priori* estimates for semilinear problems, and then describe goal-oriented variations of the standard approach to AFEM (GOAFEM). Following the recent approach of Mommer-Stevenson and Holst-Pollock for linear problems, we first establish a contraction result for the primal problem. We then develop some additional estimates that make it possible to establish contraction of the combined primal-dual quasi-error, and subsequently show convergence in the sense of the quantity of interest. Numerical experiments support the theoretical results.

1. INTRODUCTION

In this article we develop convergence theory for a class of goal-oriented adaptive finite element methods for second order semilinear equations. In particular, we establish strong contraction results for a method of this type for the problem:

$$\begin{cases} \mathcal{N}(u) := -\nabla\cdot(A\nabla u) + b(u) = f, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases} \tag{1.1}$$

with $f \in L_2(\Omega)$ and $\Omega \subset \mathbb{R}^d$ ($d = 2$ or 3) a polyhedral domain. We consider the problem with $A\colon \Omega \to \mathbb{R}^{d\times d}$ Lipschitz and almost-everywhere (a.e.) symmetric positive definite (SPD). The standard weak formulation of the primal problem reads: Find $u \in H_0^1(\Omega)$ such that

$$\langle \mathcal{N}(u), v\rangle := a(u,v) + \langle b(u), v\rangle = f(v), \quad \forall v \in H_0^1(\Omega), \tag{1.2}$$

where

$$a(u,v) = \int_\Omega A\nabla u\cdot\nabla v\, dx. \tag{1.3}$$

In many practical applications, one is more interested in certain physical quantities of the solution, referred to as "quantities of interest", such as (weighted) averages, flow rates or velocities. These quantities of interest are usually charactered by the value $g(u)$, where u is the solution and g is a linear functional associated with a particular "goal". Given a numerical approximation u_h to the solution u, goal-oriented error estimates use

---

Date: February 25, 2013.

2010 *Mathematics Subject Classification.* 65N30, 65N50, 35J61.

Key words and phrases. Adaptive methods, equations, semilinear elliptic problems, goal-oriented, quasi-orthogonality, residual-based error estimator, convergence, contraction, *a posteriori* estimates .

MH was supported in part by NSF Awards 0715146 and 0915220, and by DOD/DTRA Award HDTRA-09-1-0036.

SP was supported in part by NSF Award 0715146.

YZ was supported in part by NSF Award 0715146 and by DOD/DTRA Award HDTRA-09-1-0036.

duality techniques rather than the energy norm alone to estimate the error in the quantity of interest . The solution of the dual problem can be interpreted as the generalized Green's function, or the *influence function* with respect to the linear functional, which often quantifies the stability properties of the computed solution. There has been a lot of recent work on developing reliable and accurate *a posteriori* error estimators for goal-oriented adaptivity, see [12, 4, 5, 14, 35, 13, 18, 19, 30] and the references cited therein.

Our interest in this paper is developing a goal-oriented adaptive algorithm for semilinear problems (1.2) along with a corresponding strong contraction result, following the recent approach in [32, 23] for linear problems. One of the main challenges in the nonlinear problem that we don't see in the linear case is the dependence of the dual problem on the primal solution u. As it is not practical to work with a dual problem we cannot accurately form, we develop a method for semilinear problems in which adaptive mesh refinement is driven both by residual-based approximation to the error in u, and in a sequence of *approximate dual problems* which only depend on the numerical solution from the previous step. While globally reducing the error in the primal problem necessarily yields a good approximation to the goal error $g(u - u_h)$, methods of the type we describe here bias the error reduction in the direction of the goal-function g in the interest of achieving an accurate approximation in fewer adaptive iterations.

Contraction of the adaptive finite element algorithm for the (primal) semilinear problem (1.2) has been established in [25] and [22]. Here we recall the contraction argument for the primal problem and use a generalization of this technique to establish the contraction of a linear combination of the primal and limiting dual quasi-errors by means of a computable sequence of approximate dual problems. We relate this result to a bound on the error in the quantity of interest. Following [25], the contraction argument follows from first establishing three preliminary results for two successive AFEM approximations u_1 and u_2, and respectively $\hat{z}_1$ and $\hat{z}_2$ of the primal and limiting dual problems.

1) Quasi-orthogonality: There exists $\Lambda_G > 1$ such that
$$|||u - u_2|||^2 \le \Lambda_G |||u - u_1|||^2 - |||u_2 - u_1|||^2.$$
2) Error estimator as upper bound on error: There exists $C_1 > 0$ such that
$$|||u - u_k|||^2 \le C_1 \eta_k^2(u_k, \mathcal{T}_k), \quad k = 1, 2.$$
3) Estimator reduction: For $\mathcal{M}$ the marked set that takes refinement $\mathcal{T}_1 \to \mathcal{T}_2$, for positive constants $\lambda < 1$ and Λ_1 and any $\delta > 0$
$$\eta_2^2(v_2, \mathcal{T}_2) \le (1 + \delta)\{\eta_1^2(v_1, \mathcal{T}_1) - \lambda \eta_1^2(v_1, \mathcal{M})\} + (1 + \delta^{-1})\Lambda_1 \eta_0^2 |||v_2 - v_1|||.$$

For the primal problem, the mesh at each iteration may be marked for refinement with respect to the error indicators following the Dörfler marking strategy (cf. [11]). In the case of the dual, the limiting estimator as used in the contraction argument is related to a computable quantity. This quantity is the dual estimator, based on the residual of the approximate dual sequence. The mesh is marked for refinement with respect to this set of error indicators, which correspond to the approximate dual problem at each iteration. The transformation between limiting and approximate dual estimators couples the contraction of error in the limiting dual to the primal problem. The final result is the contraction of what we refer to here as the *combined quasi-error*
$$\bar{Q}^2(u_j, \hat{z}_j) := |||\hat{z} - \hat{z}_j|||^2 + \gamma \zeta_2^2(\hat{z}_j) + \pi |||u - u_j|||^2 + \pi \gamma_p \eta_2^2(u_j),$$
which is the sum of the quasi-error as in [8] for the limiting dual problem and a multiple of the quasi-error for the primal problem. The contraction of this property as shown in

Theorem 5.9 establishes the contraction of the error in the goal function as shown in Corollary 5.10.

Our analysis is based on the recent contraction framework for semilinear and more general nonlinear problems developed by Holst, Tsogtgerel, and Zhu [25], and by Bank, Holst, Szypowski and Zhu [2], and those for linear problems developed by Cascon, Kreuzer, Nochetto and Siebert [8], and by Nochetto, Siebert, and Veeser [34]. In addressing the goal-oriented problem we base our framework on that of Mommer and Stevenson [32] for symmetric linear problems and Holst and Pollock [23] for nonsymmetric problems. We note also two other recent convergence results in the literature for goal-oriented adaptive methods applied to self-adjoint linear problems, namely [10] and [33], both providing convergence rates in agreement with those in [32]. The analysis of the goal-oriented method for nonlinear problems is signficantly more complex than the previous analysis for linear problems in [32, 23]. Here, we are faced with analyzing linearized and approximate dual sequences as opposed to a single dual problem in order to establish contraction with respect to the quantity of interest. However, this approach allows us to establish a contraction result for the goal-oriented method, which appears to be the first result of this type for nonlinear problems. The linearized dual in the context of goal-oriented adaptive methods is described below, following e.g. Estep et. al in [14] and [15].

Outline of the paper. The remainder of the paper is structured as follows. In §2, we introduce the approximate, linearized and limiting dual problems. We briefly discuss the problem class and review some standard facts concerning conforming finite element discretization and error-estimate-driven adaptive finite element methods (AFEM). In §2.2 we include a brief summary of *a priori* estimates for the semilinear problem. In §3, we describe a goal-oriented variation of the standard approach to AFEM (GOAFEM). In §4 we discuss contraction theorems for the primal problem. In §5 we introduce additional estimates necessary for the contraction of the combined quasi-error and convergence in the sense of the quantity of interest. Lastly, in §6 we present some numerical experiments that support our theoretical results.

2. PRELIMINARIES

In this section, we state both the (nonlinear) primal problem and its finite element discretization. We then introduce the linearized dual problem, and consider some variants of this problem which are of use in the subsequent computation and analysis.

Consider the semilinear problem (1.2), where as in (1.3) we define the bilinear form

$$a(u, v) = (A\nabla u, \nabla v)$$

with $(\cdot, \cdot)$ denoting the L_2 inner-product over $\Omega \subset \mathbb{R}^d$. We make the following assumptions on the data:

Assumption 2.1 (Problem data)**.** *The problem data* $\mathbf{D} = (A, b, f)$ *satisfies*

1) $A : \Omega \to \mathbb{R}^{d\times d}$ *is Lipschitz continuous and a.e. symmetric positive-definite with*

$$\textit{ess inf}_{x\in\Omega}\lambda_{\textit{min}}(A(x)) = \mu_0 > 0,$$
$$\textit{ess sup}_{x\in\Omega}\lambda_{\textit{max}}(A(x)) = \mu_1 < \infty.$$

2) $b : \Omega \times \mathbb{R} \to \mathbb{R}$ *is smooth on the second argument. Here and in the remainder of the paper, we write* $b(u)$ *instead of* $b(x, u)$ *for simplicity. Moreover, we assume that* b *is monotone (increasing):*

$$b'(\xi) \geq 0, \ \textit{ for all } \xi \in \mathbb{R}.$$

3) $f \in L_2(\Omega)$.

The native norm is the Sobolev H^1 norm given by $\|v\|_{H^1}^2 = (\nabla v, \nabla v) + (v, v)$. Continuity of $a(\,\cdot\,,\,\cdot\,)$ follows from the Hölder inequality, and bounding the L_2 norm of the function and its gradient by the H^1 norm

$$a(u, v) \le \mu_1 \|u\|_{H^1}\|v\|_{H^1} = M_{\mathcal{E}}\|u\|_{H^1}\|v\|_{H^1}. \tag{2.1}$$

Define the energy semi-norm by the principal part of the differential operator $|\!|\!| v |\!|\!|^2 := a(v, v)$. The coercivity of $a(\,\cdot\,,\,\cdot\,)$ follows from the Poincaré inequality with constant C_Ω

$$a(v, v) \ge \mu_0 |v|_{H^1}^2 \ge C_\Omega \mu_0 \|v\|_{H^1}^2 = m_{\mathcal{E}}^2 \|v\|_{H^1}^2, \tag{2.2}$$

which establishes the energy semi-norm as a norm. Putting this together with (2.1) establishes the equivalence between the native and energy norms.

2.1. **Linearized dual problems.** Given a linear funcitonal $g \in L_2(\Omega)$, the objective in goal-oriented error estimation is to relate the residual to the error in the quantity of interest. This involves solving a dual problem whose solution z satisfies the relation $g(u - u_h) = \langle R(u_h), z\rangle$. In the linear case, the appropriate dual problem is the formal adjoint of the primal (cf. [31, 23]). For b nonlinear, the primal problem (1.2) does not have an exact formal adjoint. In this case we obtain the dual by linearization. Formally, given a numerical approximation u_j to the exact solution u, the residual is given by

$$R(u_j) := f - \mathcal{N}(u_j) = \mathcal{N}(u) - \mathcal{N}(u_j).$$

If $z^j \in H_0^1(\Omega)$ solves the following linearized dual problem

$$a(z^j, v) + \langle \mathcal{B}_j z^j, v\rangle = g(v), \quad \forall v \in H_0^1(\Omega), \tag{2.3}$$

where $g(v) := \int_\Omega g v dx$ and the operator $\mathcal{B}_j$ is given by

$$\mathcal{B}_j := \int_0^1 b'(\xi u + (1 - \xi)u_j)\, d\xi = \int_0^1 b'(u_j + (u - u_j)\xi)\, d\xi, \tag{2.4}$$

then the goal-oriented error $g(e_j)$ can be represented exactly by the inner product of z^j and $R(u_j)$:

$$g(e_j) = \langle R(u_j), z^j\rangle.$$

In fact, by definition of the residual $R(u_j)$, we have

$$\langle z^j, R(u_j)\rangle = a(z^j, e_j) + \langle z^j, b(u) - b(u_j)\rangle = a(z^j, e_j) + \langle \mathcal{B}_j z^j, v\rangle = g(e_j).$$

Here we used the integral mean value identity:

$$b(u) - b(u_j) = \int_0^1 b'(u_j + (u - u_j)\xi)\, d\xi (u - u_j) = \mathcal{B}_j(u - u_j).$$

The derivation and numerical use of the linearized dual problem is further discussed in [15, 14, 21].

Unfortunately, the dual problem (2.3) is not practical because the operator $\mathcal{B}_j$ depends on the exact solution u. In order to define a computable dual operator, we introduce the approximate operator $b'(u_j)$, which lead to the following approximate dual problem: Find $\hat{z}^j \in H_0^1(\Omega)$ such that

$$a(\hat{z}^j, v) + \langle b'(u_j)\hat{z}^j, v\rangle = g(v), \quad \forall v \in H_0^1(\Omega). \tag{2.5}$$

The equation (2.5) is instrumental for defining a computable *a posteriori* error indicator for the dual problem.

A further difficulty arises in the analysis of the goal-oriented adaptive algorithm driven by the *a posteriori* error estimators for the approximate dual problem (2.5). Due to the dependence on u_j, (2.5) changes at each step of the adaptive algorithm. This is one of the essential differences of the nonlinear problem as compared to the linear cases in the previous literature (cf. [32, 23]). To handle this obstacle, we introduce the limiting dual problem: Find $\hat{z} \in H_0^1(\Omega)$ such that

$$a(\hat{z}, v) + \langle b'(u)\hat{z}, v\rangle = g(v), \quad \forall v \in H_0^1(\Omega). \tag{2.6}$$

While the operator $b'(u)$ is a function of the exact solution u and is not a computable quantity, it is the operator used in the limit of both the linearized dual (2.3) and approximate dual problems (2.5) as $u_j \to u$. Therefore, both the linearized and approximate sequences approach the same limiting problem (2.6). Our contraction result in Theorem 5.9 is written with respect to the limiting dual problem as defined by the operator $b'(u)$.

2.2. **Finite Element Approximation.** For a given conforming, shape-regular triangulation $\mathcal{T}$ of Ω consisting of close simplices $T \in \mathcal{T}$, we define the finite element space

$$\mathbb{V}_{\mathcal{T}} := H_0^1(\Omega) \cap \prod_{T\in\mathcal{T}} \mathbb{P}_n(T) \quad \text{and } \mathbb{V}_k := \mathbb{V}_{\mathcal{T}_k}, \tag{2.7}$$

where $\mathbb{P}_n(T)$ is the space of polynomials degree degree n over T. For any subset $\mathcal{S} \subseteq \mathcal{T}$,

$$\mathbb{V}_{\mathcal{T}}(\mathcal{S}) := H_0^1(\Omega) \cap \prod_{T\in\mathcal{S}} \mathbb{P}_n(T). \tag{2.8}$$

Given a triangulation $\mathcal{T}$, we denote $h_{\mathcal{T}} := \max_{T\in\mathcal{T}} h_T$ where $h_T := |T|^{1/d}$. In particular, we denote $h_0 := h_{\mathcal{T}_0}$ for an initial (conforming, shape-regular) triangulation $\mathcal{T}_0$ of Ω. Then the adaptive algorithm discussed below generates a nested sequence of conforming refinements $\{\mathcal{T}_k\}$, with $\mathcal{T}_k \geq \mathcal{T}_j$ for $k \geq j$ meaning that $\mathcal{T}_k$ is a conforming triangulation of Ω based on certain refinements of $\mathcal{T}_j$. With this notation, we also simply denote by $\mathbb{V}_k := \mathbb{V}_{\mathcal{T}_k}$ the finite element space defined on $\mathcal{T}_k$.

The finite element approximation of the primal problem (1.2) reads: Find $u_k \in \mathbb{V}_k$ such that

$$a(u_k, v_k) + \langle b(u_k), v_k\rangle = f(v_k), \; v_k \in \mathbb{V}_k, \tag{2.9}$$

and the finite element approximation of (2.5) linearized about u_j is given by: Find $\hat{z}_k^j \in \mathbb{V}_k$ such that

$$a(\hat{z}_k^j, v_k) + \langle b'(u_j)\hat{z}_k^j, v_k\rangle = g(v_k) \quad \text{for all } v_k \in \mathbb{V}_k. \tag{2.10}$$

Finally, for the purpose of analysis, we require the discrete limiting dual problem (cf. (2.6)) given by: Find $\hat{z}_k \in \mathbb{V}_k$ such that

$$a(\hat{z}_k, v_k) + \langle b'(u)\hat{z}_k, v_k\rangle = g(v_k) \quad \text{for all } v_k \in \mathbb{V}_k. \tag{2.11}$$

Existence and uniqueness of solutions to the primal problems (1.2) and (2.9) follow from standard variational or fixed-point arguments as in [38] and [29]. For the dual problems (2.5)-(2.6) and (2.10)-(2.11) the existence and uniqueness of solutions follow from the standard Lax-Milgram Theorem as in [17], since we assumed that $b'(\xi) \geq 0$.

We make the following assumption on the a priori L_∞ bounds of the solutions to the primal problems (1.2) and (2.9):

Assumption 2.2 (*A priori bounds*)**.** *Let u and u_k be the solution to* (1.2) *and* (2.9)*,respectively. We assume that there are $u_-, u_+ \in L_\infty$ which satisfy*

$$u_-(x) \leq u(x), u_k(x) \leq u_+(x) \text{ for almost every } x \in \Omega. \tag{2.12}$$

Remark 2.3. *The L_∞ bound on u follows from the standard maximum principle, as discussed in* [3, Theorem 2.4] *and* [24, Theorem 2.3]. *There has been a lot of literature on the L_∞ bounds on the discrete solution, usually with additional angle condition of the triangulation (cf.* [28, 26, 27, 24] *and the references cited therein). On the other hand, if b satisfies the (sub)critical growth condition, as stated in* [3, Assumption (A4)], *then the L_∞ bounds on the discrete solution u_k are satisfied without angle conditions on the mesh, see* [3] *for more details.*

Assumption 2.1 together with Assumption 2.2 yield the following properties on the continuous and discrete solutions as summarized below.

Proposition 2.4. *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. The following properties hold:*

1) *b is Lipschitz on $[u_-, u_+] \cap H_0^1(\Omega)$ for a.e. $x \in \Omega$ with constant B.*
2) *b' is Lipschitz on $[u_-, u_+] \cap H_0^1(\Omega)$ for a.e. $x \in \Omega$ with constant Θ.*
3) *Let $\hat{z}$ the solution to (2.6), $\hat{z}_j^j$ the solution to (2.10) and $\hat{z}_j$ the solution to (2.11). Then there are $z_-, z_+ \in L_\infty$ which satisfy*

$$z_-(x) < \hat{z}(x), \hat{z}_j(x), \hat{z}_j^j(x) \le z_+(x) \text{ for almost every } x \in \Omega,\ j \in \mathbb{N}. \tag{2.13}$$

3. GOAL ORIENTED AFEM

In this section, we describe the goal oriented adaptive finite element method (GOAFEM), which is based on the standard AFEM algorithm:

$$\text{SOLVE} \rightarrow \text{ESTIMATE} \rightarrow \text{MARK} \rightarrow \text{REFINE}\,.$$

Below, we explain each procedure.

Procedure SOLVE. The procedure SOLVE involves solving (2.9) for u_j, computing $b'(u_j)$ to form problem (2.10) and solving (2.10) for $\hat{z}_j^j$. In the analysis that follows, we assume for simplicity that the exact Galerkin solution is found on each mesh refinement. In practice the nonlinear problem (2.9) may be solved by a standard inexact Newton + multilevel algorithm as in [2]. The approximate dual problem (2.10) may be solved by any standard linear-time iterative method.

Procedure ESTIMATE. We use a fairly standard residual-based element-wise error estimator for both primal and approximate dual problems. Recall that the residual of the primal problem is given by $R(v) = f - \mathcal{N}(v)$ with $\mathcal{N}(v) = -\nabla \cdot (A\nabla v) + b(v)$. For the limiting and approximate dual problems, we define the local strong form by $\hat{\mathcal{L}}^*(v) := -\nabla \cdot (A\nabla v) + b'(u)(v)$, and $\hat{\mathcal{L}}_j^*(v) := -\nabla \cdot (A\nabla v) + b'(u_j)(v)$. The limiting and approximate dual residuals given respectively by

$$R^*(v) := g - \hat{\mathcal{L}}^*(v), \text{ and } \hat{R}_j^*(v) := g - \hat{\mathcal{L}}_j^*(v). \tag{3.1}$$

The *jump residual* for both the primal and linearized dual problems is:

$$J_T(v) := [\![[A\nabla v] \cdot n]\!]_{\partial T},$$

where $[\![\,\cdot\,]\!]$ is given by $[\![\phi]\!]_{\partial T} := \lim_{t\to 0} \phi(x + tn) - \phi(x - tn)$and n is taken to be the appropriate outward normal defined on ∂T. On boundary edges σ_b we have $[\![[A\nabla v] \cdot n]\!]_{\sigma_b} \equiv 0$ so that $[\![[A\nabla v] \cdot n]\!]_{\partial T} = [\![[A\nabla v] \cdot n]\!]_{\partial T \cap \Omega}$. The error indicator for the primal problem (2.9) is given by

$$\eta_{\mathcal{T}}^2(v, T) := h_T^2 \|R(v)\|_{L_2(T)}^2 + h_T \|J_T(v)\|_{L_2(\partial T)}^2, \quad v \in \mathbb{V}_{\mathcal{T}}. \tag{3.2}$$

Similarly, the dual error-indicator is given by the approximate residual

$$\zeta^2_{\mathcal{T},j}(w,T) \coloneqq h_T^2\|\hat{R}^*_j(w)\|^2_{L_2(T)} + h_T\|J_T(w)\|^2_{L_2(\partial T)}, \quad w \in \mathbb{V}_{\mathcal{T}}. \tag{3.3}$$

This dual indicator is defined in terms of the approximate dual operator $b'(u_j)$ as this is a computable quantity given an an approximation u_j. In addition, for purpose of analysis we define the limiting dual error-indicator by

$$\zeta^2_{\mathcal{T}}(w,T) \coloneqq h_T^2\|\hat{R}^*(w)\|^2_{L_2(T)} + h_T\|J_T(w)\|^2_{L_2(\partial T)}, \quad w \in \mathbb{V}_{\mathcal{T}}. \tag{3.4}$$

We remark that the limiting dual indicator as given by (3.4) is not computable. For any given subset $\mathcal{S} \subset \mathcal{T}$, the error estimators on $\mathcal{S}$ are given by the l_2 sum of error indicators over elements in the space.

$$\eta^2_{\mathcal{T}}(v,\mathcal{S}) \coloneqq \sum_{T\in\mathcal{S}} \eta^2_{\mathcal{T}}(v,T), \quad v \in \mathbb{V}_{\mathcal{T}}.$$

The dual energy estimator is:

$$\zeta^2_{\mathcal{T},j}(w,\mathcal{S}) \coloneqq \sum_{T\in\mathcal{S}} \zeta^2_{\mathcal{T},j}(w,T), \quad w \in \mathbb{V}_{\mathcal{T}},$$

and the limiting estimator

$$\zeta^2_{\mathcal{T}}(w,\mathcal{S}) \coloneqq \sum_{T\in\mathcal{S}} \zeta^2_{\mathcal{T}}(w,T), \quad w \in \mathbb{V}_{\mathcal{T}}.$$

To simplify the notation, below we will omit "$\mathcal{S}$" in the above definitions if $\mathcal{S} = \mathcal{T}$ and we will use η_k to denote $\eta_{\mathcal{T}_k}$, and similarly use $\zeta_{k,\cdot}$ to denote $\zeta_{\mathcal{T}_k,\cdot}$.

As in [8] it is not difficult to verify that the indicators for the primal and approximate (respectively limiting) dual problems satisfy the monotonicity property for $v \in \mathbb{V}(\mathcal{T}_1)$ and $\mathcal{T}_2 \geq \mathcal{T}_1$

$$\eta_2(v,\mathcal{T}_2) \leq \eta_1(v,\mathcal{T}_1),\ \zeta_{2,j}(v,\mathcal{T}_2) \leq \zeta_{1,j}(v,\mathcal{T}_1) \ \text{ and } \ \zeta_2(v,\mathcal{T}_2) \leq \zeta_1(v,\mathcal{T}_1). \tag{3.5}$$

For an element $T \in \mathcal{T}_2 \cap \mathcal{T}_1$

$$\eta_2(v,T) = \eta_1(v,T),\ \zeta_{2,j}(v,T) = \zeta_{1,j}(v,T) \ \text{ and } \ \zeta_2(v,T) = \zeta_1(v,T). \tag{3.6}$$

Procedure MARK. The Dörfler marking strategy for the goal-oriented problem is based on the following steps as in [32]:

1) Given $\theta \in (0,1)$, mark sets for each of the primal and dual problems:
   - Mark a set $\mathcal{M}_p \subset \mathcal{T}_k$ such that
   $$\eta^2_k(u_k,\mathcal{M}_p) \geq \theta^2\eta^2_k(u_k,\mathcal{T}_k). \tag{3.7}$$
   - Mark a set $\mathcal{M}_d \subset \mathcal{T}_k$ such that
   $$\zeta^2_{k,k}(\hat{z}^k_k,\mathcal{M}_d) \geq \theta^2\zeta^2_{k,k}(\hat{z}^k_k,\mathcal{T}_k). \tag{3.8}$$
2) Let $\mathcal{M} = \mathcal{M}_p \cup \mathcal{M}_d$ the union of sets found for the primal and dual problems respectively.

As in [23] the set $\mathcal{M}$ differs from that in [32], where the set of lesser cardinality between $\mathcal{M}_p$ and $\mathcal{M}_d$ is used. As seen in (3.8) the mesh is marked with respect to the dual indicators of the approximate-sequence solutions $\hat{z}^k_k$ as these are computable quantities. Sets $\mathcal{M}_p$ and $\mathcal{M}_d$ with optimal cardinality (up to a factor of 2) can be chosen in linear time by binning the elements rather than performing a full sort [32].

Procedure REFINE. The refinement (including the completion) is performed according to newest vertex bisection which was first proposed in [36]. It has been proved that the bisection procedure will preserve the shape-regularity of the initial triangulation $\mathcal{T}_0$. The complexity and other properties of this procedure are now well-understood (see for example [6] and the references cited therein), and will simply be exploited here.

4. CONTRACTION FOR THE PRIMAL PROBLEM

Here we discuss the contraction of the primal problem (1.2), recalling results from [25], [24] and [2]. The contraction argument relies on three main convergence results, namely quasi-orthogonality, error-estimator as upper bound on error and estimator reduction. We include the analogous results here for the limiting dual problem when they are identical or nearly identical.

4.1. Quasi-orthogonality.

Orthogonality in the energy-norm $|||u-u_2|||^2 = |||u-u_1|||^2 - |||u_2-u_1|||^2$ does not generally hold in the semilinear problem. We rely on the weaker quasi-orthogonality result to establish contraction of AFEM (GOAFEM). The proof of the quasi-orthogonality relies on the following L_2-lifting property.

Lemma 4.1 (L_2-lifting)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let u be the exact solution to (1.2), and $u_1 \in \mathbb{V}_1$ the Galerkin solution to (2.9). Let $w \in H^{1+s}(\Omega) \cap H^1_0(\Omega)$ for some $0 < s \leq 1$ be the solution to the dual problem: Find $w \in H^1_0(\Omega)$ such that*

$$a(w,v) + \langle \mathcal{B}_1 w, v\rangle = \langle u - u_1, v\rangle, \quad v \in H^1_0(\Omega), \tag{4.1}$$

where the operator $\mathcal{B}_1$ is defined by $\mathcal{B}_1 := \int_0^1 b'(\xi u + (1-\xi)u_1)\, d\xi$. As in [9, 16, 1] we assume the regularity

$$|w|_{H^{1+s}(\Omega)} \leq K_R \|u - u_1\|_{L_2(\Omega)} \tag{4.2}$$

based on the continuity of the diffusion coefficients a_{ij} and $\mathcal{B}_1 \in L_\infty(\Omega)$. Then

$$\|u-u_1\|_{L_2} \leq C_* h_0^s |||u-u_1|||. \tag{4.3}$$

Proof. The proof follows the standard duality arguments in [1], [23] and [7], adapted for the semilinear problem. Let $\mathcal{I}^h : H^1_0(\Omega) \to \mathbb{V}_1$ be a quasi-interpolator, satisfying

$$\|w - \mathcal{I}^h w\|_{H^1} \leq C_{\mathcal{I}} h^s_{\mathcal{T}_1} |w|_{H^{1+s}} \tag{4.4}$$

$$\|w - \mathcal{I}^h w\|_{L_2} \leq \hat{C}_{\mathcal{I}} h^{1+s}_{\mathcal{T}_1} |w|_{H^{1+s}}. \tag{4.5}$$

as discussed in [1], [37] and [23].

Consider the linearized dual problem (4.1) with $v = u - u_1 \in H^1_0(\Omega)$ expressed in primal form

$$a(u-u_1, w) + \langle \mathcal{B}_1(u-u_1), w\rangle = \|u-u_1\|^2_{L_2}. \tag{4.6}$$

By Galerkin orthogonality, for $\mathcal{I}^h w \in \mathbb{V}_1$

$$a(u-u_1, \mathcal{I}^h w) + \langle \mathcal{B}_1(u-u_1), \mathcal{I}^h w\rangle = 0. \tag{4.7}$$

Subtracting (4.7) from (4.6)

$$a(u-u_1, w-\mathcal{I}^h w) + \langle b(u) - b(u_1), w - \mathcal{I}^h w\rangle = \|u-u_1\|^2_{L_2}. \tag{4.8}$$

Then by (2.1) continuity of $a(\,\cdot\,,\,\cdot\,)$, the Hölder inequality and Lipschitz continuity of b (Proposition 2.4):

$$\|u-u_1\|^2_{L_2} \leq M_{\mathcal{E}} \|u-u_1\|_{H^1} \|w - \mathcal{I}^h w\|_{H^1} + B\|u-u_1\|_{L_2} \|w-\mathcal{I}^h w\|_{L_2}. \tag{4.9}$$

By coercivity (2.2), interpolation estimate (4.4), and regularity (4.2) on the first term on the RHS of (4.9)

$$\begin{aligned}M_{\mathcal{E}}\|u-u_1\|_{H^1}\|w-\mathcal{I}^h w\|_{H^1} &\le \frac{M_{\mathcal{E}}}{m_{\mathcal{E}}}C_{\mathcal{I}}h_0^s|\!|\!|u-u_1|\!|\!||w|_{H^{1+s}}\\ &\le \frac{M_{\mathcal{E}}}{m_{\mathcal{E}}}K_R C_{\mathcal{I}}h_0^s|\!|\!|u-u_1|\!|\!|\|u-u_1\|_{L_2}. \end{aligned}\tag{4.10}$$

For the second term of (4.9), apply (4.5) followed by (4.2) and coercivity to the interpolation error yielding

$$\begin{aligned}B\|u-u_1\|_{L_2}\|w-\mathcal{I}^h w\|_{L_2} &\le B\hat{C}_{\mathcal{I}}h_0^{1+s}\|u-u_1\|_{L_2}|w|_{H^{1+s}}\\ &\le K_R B\hat{C}_{\mathcal{I}}h_0^{1+s}\|u-u_1\|_{L_2}\|u-u_1\|_{L_2}\\ &\le (m_{\mathcal{E}}^{-1}K_R B\hat{C}_{\mathcal{I}}h_0)h_0^s\|u-u_1\|_{L_2}|\!|\!|u-u_1|\!|\!|. \end{aligned}\tag{4.11}$$

Applying (4.10) and (4.11) to (4.9), we obtain

$$\|u-u_1\|_{L_2} \le m_{\mathcal{E}}^{-1}K_R\left(M_{\mathcal{E}}C_{\mathcal{I}}+B\hat{C}_{\mathcal{I}}h_0\right)h_0^s|\!|\!|u-u_1|\!|\!|. \tag{4.12}$$

This completes the proof. □

Remark 4.2. *This proof yields the related result for $u_i \in \mathbb{V}_i$ the solution to (2.9), $i=1,2$*

$$\|u_2-u_1\|_{L_2} \le C_* h_0^s |\!|\!|u_2-u_1|\!|\!|. \tag{4.13}$$

The proof of (4.13) follows by replacing u by u_2 in 4.1. In particular, the dual problem (4.1) is replaced by: Find $w \in \mathbb{V}_2$ such that

$$a(w,v)+\langle \mathcal{B}_{12}w,v\rangle = (u_2-u_1,v), \quad v\in\mathbb{V}_2, \tag{4.14}$$

where the operator $\mathcal{B}_{12} := \int_0^1 b'(\xi u_2 + (1-\xi)u_1)\,d\xi$.

Remark 4.3. *As the dual problem (4.1) changes at each iteration, so may the regularity constant as given by (4.2) as well as the interpolation constants as given by (4.4) and (4.5). As such, the previous lemma shows a $C_{*,k}$ for $k=1,2,\ldots$. As the algorithm is run finitely many times, we consolidate these $C_{*,k}$ into a single constant C_* for simplicity of presentation.*

Now we are in position to show the quasi-orthogonality.

Lemma 4.4 (Quasi-orthogonality)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let $\mathcal{T}_1, \mathcal{T}_2$ be two conforming triangulation of Ω with $\mathcal{T}_2 \ge \mathcal{T}_1$. Let $u \in H_0^1(\Omega)$ the exact solution to (1.2), $u_i \in \mathbb{V}_i$ the solution to (2.9), $i=1,2$. There exists a constant $C_*>0$ depending on the problem data $\mathbf{D}$ and initial mesh $\mathcal{T}_0$, and a number $0<s\le 1$ related to the angles of $\partial\Omega$, such that if the meshsize h_0 of the initial mesh satisfies $\bar{\Lambda} := Bm_{\mathcal{E}}^{-1}C_*h_0^s < 1$, then*

$$|\!|\!|u-u_2|\!|\!|^2 \le \Lambda|\!|\!|u-\bar{v}|\!|\!|^2 - |\!|\!|u_2-\bar{v}|\!|\!|^2, \quad \forall \bar{v}\in\mathbb{V}_2, \tag{4.15}$$

and in particular for $\bar{v}=u_1\in\mathbb{V}_1\subset\mathbb{V}_2$

$$|\!|\!|u-u_2|\!|\!|^2 \le \Lambda_G|\!|\!|u-u_1|\!|\!|^2 - |\!|\!|u_2-u_1|\!|\!|^2, \tag{4.16}$$

where

$$\Lambda := (1-Bm_{\mathcal{E}}^{-1}C_*h_0^s)^{-1} \ \textit{and}\ \ \Lambda_G := (1-BC_*^2h_0^{2s})^{-1}$$

and C_* is the constant from Lemma 4.1.

Proof. For any given $\bar{v} \in \mathbb{V}_2$, we have

$$|\!|\!| u - u_2 |\!|\!|^2 = |\!|\!| u - \bar{v} |\!|\!|^2 - |\!|\!| \bar{v} - u_2 |\!|\!|^2 + 2a(u - u_2, \bar{v} - u_2). \tag{4.17}$$

By Galerkin orthogonality

$$a(u - u_2, v) + \langle b(u) - b(u_2), v \rangle = 0 \text{ for all } v \in \mathbb{V}_2, \tag{4.18}$$

and taking $v = \bar{v} - u_2$ in (4.18), we have

$$\begin{aligned} 2a(u - u_2, \bar{v} - u_2) &\le 2|\langle b(u) - b(u_2), \bar{v} - u_2 \rangle| \\ &\le 2B \|u - u_2\|_{L_2} \|\bar{v} - u_2\|_{L_2}. \end{aligned} \tag{4.19}$$

Here we used Hölder inequality and the Lipschitz property on b (cf. Proposition 2.4).

In the case of (4.15) applying L_2-lifting Lemma 4.1 to the first factor on the RHS and (2.2) coercivity to the second followed by Young's inequality, we obtain

$$\begin{aligned} 2B\|u - u_2\|_{L_2} \|\bar{v} - u_2\|_{L_2} &\le 2Bm_{\mathcal{E}}^{-1} C_* h_0^s |\!|\!| u - u_2 |\!|\!| \, |\!|\!| \bar{v} - u_2 |\!|\!| \\ &\le Bm_{\mathcal{E}}^{-1} C_* h_0^s |\!|\!| u - u_2 |\!|\!|^2 + Bm_{\mathcal{E}}^{-1} C_* h_0^s |\!|\!| \bar{v} - u_2 |\!|\!|^2. \end{aligned} \tag{4.20}$$

Applying (4.20) via (4.19) to (4.17)

$$(1 - Bm_{\mathcal{E}}^{-1} C_* h_0^s) |\!|\!| u - u_2 |\!|\!|^2 \le |\!|\!| u - \bar{v} |\!|\!|^2 - (1 - Bm_{\mathcal{E}}^{-1} C_* h_0^s) |\!|\!| \bar{v} - u_2 |\!|\!|^2.$$

Assuming $\bar{\Lambda} := Bm_{\mathcal{E}}^{-1} C_* h_0^s < 1$, we have

$$|\!|\!| u - u_2 |\!|\!|^2 \le \Lambda |\!|\!| u - \bar{v} |\!|\!|^2 - |\!|\!| \bar{v} - u_2 |\!|\!|^2 \tag{4.21}$$

with $\Lambda = (1 - Bm_{\mathcal{E}}^{-1} C_* h_0^s)^{-1}$.

In the case of (4.16) applying L_2-lifting 4.1 to each norm on the RHS of (4.19) by means of Remark 4.2 then applying Young's inequality

$$\begin{aligned} 2B\|u - u_2\|_{L_2} \|u_1 - u_2\|_{L_2} &\le 2Bh_0^{2s} C_*^2 |\!|\!| u - u_2 |\!|\!| \, |\!|\!| u_1 - u_2 |\!|\!| \\ &\le Bh_0^{2s} C_*^2 |\!|\!| u - u_2 |\!|\!|^2 + BC_*^2 h_0^{2s} |\!|\!| u_1 - u_2 |\!|\!|^2. \end{aligned} \tag{4.22}$$

Following the same procedure as above yields

$$|\!|\!| u - u_2 |\!|\!|^2 \le \Lambda_G |\!|\!| u - u_1 |\!|\!|^2 - |\!|\!| u_1 - u_2 |\!|\!|^2 \tag{4.23}$$

with $\Lambda_G = (1 - BC_*^2 h_0^{2s})^{-1}$ with the weaker mesh assumption $\bar{\Lambda}_G := BC_*^2 h_0^{2s} < 1$. $\square$

We note that the second Galerkin orthogonality estimate (4.23) sharpens our results but is not essential to establishing them.

4.2. **Error Estimator as Global Upper-bound.**

The second key result for the contraction of the primal problem is the error estimator as a global upper bound on the energy error, up to a global constant. The result for the semilinear problem is established in [25] and [2] with a clear generalization to the approximate dual sequence, also see [8] and [31] for the linear cases. The proof of this result follows from the general *a posteriori* error estimation framework developed in [39, 40].

Lemma 4.5 (Error estimator as global upper-bound)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let $\mathcal{T}_k$ be a conforming refinement of $\mathcal{T}_0$. Let $u \in H_0^1(\Omega)$ and $u_k \in \mathbb{V}_k$ be the solutions to (1.2) and (2.9), respectively. Similarly, let $\hat{z} \in H_0^1(\Omega)$ and $\hat{z}_k \in \mathbb{V}_k$ be the solutions to (2.6) and (2.11), respectively. Then there is a global constant C_1 depending only on the problem data* $\mathbf{D}$ *and initial mesh $\mathcal{T}_0$ such that*

$$|\!|\!| u - u_k |\!|\!| \le C_1 \eta_k(u_k) \tag{4.24}$$

and

$$|\!|\!| \hat{z} - \hat{z}_k |\!|\!| \le C_1 \zeta_k(\hat{z}_k). \tag{4.25}$$

4.3. **Estimator Reduction.** The local Lipschitz property as in [25], analogous to the local perturbation property established in [8], is a key step in establishing estimator reduction leading to the contraction result. For any $T \in \mathcal{T}$, we denote

$$\omega_T := T \cup \{T' \in \mathcal{T} \,\big|\, T \cap T' \text{ is a true-hyperface of } T\}. \tag{4.26}$$

Here, for a d-simplex T, a true-hyperface is a $d-1$ sub-simplex of T, e.g., a face in 3D or an edge in 2D. We also define the data estimator on each element $T \in \mathcal{T}$ as

$$\eta_{\mathcal{T}}^2(\mathbf{D}, T) = h_T^2 \left(\|\text{div} A\|_{L_\infty(T)}^2 + h_T^{-2} \|A\|_{L_\infty(\omega_T)}^2 + B^2 \right), \tag{4.27}$$

and denote $\eta_{\mathcal{T}}(\mathbf{D}, \mathcal{S}) = \max_{T \in \mathcal{S}} \eta_{\mathcal{T}}(\mathbf{D}, T)$ for any subset $\mathcal{S} \subseteq \mathcal{T}$. In particular, we denote by $\eta_0 := \eta_{\mathcal{T}_0}(\mathbf{D}, \mathcal{T}_0)$ the data estimator on the initial mesh. As the grid is refined, the data estimator satisfies the monotonicity property for refinements $\mathcal{T}_2 \geq \mathcal{T}_1$ (cf. [8]):

$$\eta_2(\mathbf{D}, \mathcal{T}_2) \leq \eta_1(\mathbf{D}, \mathcal{T}_1). \tag{4.28}$$

Lemma 4.6 (Local Lipschitz Property)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let $\mathcal{T}$ be a conforming refinement of $\mathcal{T}_0$. Then for all $T \in \mathcal{T}$ and for any $v, w \in \mathbb{V}_{\mathcal{T}}$*

$$|\eta_{\mathcal{T}}(v, T) - \eta_{\mathcal{T}}(w, T)| \leq \bar{\Lambda}_1 \eta_{\mathcal{T}}(\mathbf{D}, T) \|v - w\|_{H^1(\omega_T)}. \tag{4.29}$$

The constant $\bar{\Lambda}_1 > 0$ depends on the dimension d and the initial mesh $\mathcal{T}_0$.

Proof. The proof follows those in [8] and [23], and we sketch the proof below. From (3.2)

$$\eta_{\mathcal{T}}^2(v, T) := h_T^2 \|R(v)\|_{L_2(T)}^2 + h_T \|J_T(v)\|_{L_2(\partial T)}^2, \quad v \in \mathbb{V}_{\mathcal{T}}. \tag{4.30}$$

Set $e = v - w$ and by definition of the residual, we get

$$\begin{aligned} R(v) &= f - \mathcal{N}(w + e) \\ &= f + \nabla \cdot (A \nabla w) - b(w) + \nabla \cdot (A \nabla e) - \int_0^1 b'(w + \xi e) \, d\xi e \\ &= R(w) + \mathcal{D}(e), \end{aligned}$$

where $\mathcal{D}(e) := \nabla \cdot (A \nabla e) - \int_0^1 b'(w + \xi e) \, d\xi e$. Using the generalized triangle-inequality

$$\sqrt{(a+b)^2 + (c+d)^2} \leq \sqrt{a^2 + c^2} + b + d, \qquad \text{for } a, b, c, d > 0$$

and linearity of the jump residual we have

$$\begin{aligned} \eta_{\mathcal{T}}(v, T) &= \left(h_T^2 \|R(w) + \mathcal{D}(e)\|_{L_2(T)}^2 + h_T \|J(w) + J(e)\|_{L_2(\partial T)}^2 \right)^{1/2} \\ &\leq \eta_{\mathcal{T}}(w, T) + h_T \|\mathcal{D}(e)\|_{L_2(T)} + h_T^{1/2} \|J(e)\|_{L_2(\partial T)}. \end{aligned} \tag{4.31}$$

For the second term of (4.31), by triangle inequality we obtain

$$\|\mathcal{D}(e)\|_{L_2(T)} \leq \|\nabla \cdot (A \nabla e)\|_{L_2(T)} + \left\| \int_0^1 b'(w + \xi e) \, d\xi e \right\|_{L_2(T)}. \tag{4.32}$$

By the inverse inequality, the diffusion term satisfies the bound

$$\begin{aligned} \|\nabla \cdot (A \nabla e)\|_{L_2(T)} &\leq \|\text{div} A \cdot \nabla e\|_{L_2(T)} + \|A : D^2 e\|_{L_2(T)} \\ &\leq \left(\|\text{div} A\|_{L_\infty(T)} + C_I h_T^{-1} \|A\|_{L_\infty(T)} \right) \|\nabla e\|_{L_2(T)}, \end{aligned} \tag{4.33}$$

where $D^2 e$ is the Hessian of e. The second term in (4.32) is bounded by

$$\left\| \int_0^1 b'(w + \xi e) \, d\xi e \right\|_{L_2(T)} \leq B \|e\|_{L_2(T)}. \tag{4.34}$$

The jump term in (4.31) satisfies

$$\begin{aligned}\|J(e)\|_{L_2(\partial T)} &\le 2(d+1)\, C_T h_T^{-1/2}\|A\|_{L_\infty(\omega_T)}\|\nabla e\|_{L_2(\omega_T)}\\ &= C_J h_T^{-1/2}\|A\|_{L_\infty(\omega_T)}\|\nabla e\|_{L_2(\omega_T)},\end{aligned} \tag{4.35}$$

where C_T depends only on the shape-regularity of the triangulation. Putting together (4.31), (4.33), (4.34) and (4.35), we obtain

$$\begin{aligned}\eta_{\mathcal{T}}(v,T) &\le \eta_{\mathcal{T}}(w,T) + h_T\left(\|\mathrm{div}A\|_{L_\infty(T)} + (C_I + C_J)h_T^{-1}\|A\|_{L_\infty(\omega_T)} + B\right)\|e\|_{H^1(\omega_T)}\\ &\le \eta_{\mathcal{T}}(w,T) + C_{TOT}\,\eta_{\mathcal{T}}(\mathbf{D},T)\|v-w\|_{H^1(\omega_T)}.\end{aligned} \tag{4.36}$$

This completes the proof. $\square$

The local perturbation property as demonstrated in Lemma 4.6 (respectively, Lemma 5.4 below) leads to estimator reduction, one of the three key ingredients for contraction of the both the primal and combined quasi-errors. This result holds for both the primal and limiting dual problems, whose proof can be found in [8, Corollary 2.4] or [23, Theorem 3.4].

Theorem 4.7 (Estimator reduction)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let $\mathcal{T}_1$ be a conforming refinements of $\mathcal{T}_0$, $\mathcal{M} \subset \mathcal{T}_1$ be the marked set, and $\mathcal{T}_2 = REFINE(\mathcal{T}_1, \mathcal{M})$. Let*

$$\Lambda_1 := (d+2)\bar{\Lambda}_1^2 m_{\mathcal{E}}^{-2} \quad \text{and} \quad \lambda := 1 - 2^{-1/d} > 0$$

with $\bar{\Lambda}_1$ from Lemma 4.6 (local Lipschitz property). Then for any $v_1 \in \mathbb{V}_1$ and $v_2 \in \mathbb{V}_2$ and $\delta > 0$

$$\eta_2^2(v_2,\mathcal{T}_2) \le (1+\delta)\left\{\eta_1^2(v_1,\mathcal{T}_1) - \lambda\eta_1^2(v_1,\mathcal{M})\right\} + (1+\delta^{-1})\Lambda_1\eta_0^2|||v_2 - v_1|||^2. \tag{4.37}$$

Analogously for the limiting dual problem

$$\zeta_2^2(v_2,\mathcal{T}_2) \le (1+\delta)\left\{\zeta_1^2(v_1,\mathcal{T}_1) - \lambda\zeta_1^2(v_1,\mathcal{M})\right\} + (1+\delta^{-1})\Lambda_1\eta_0^2|||v_2 - v_1|||^2. \tag{4.38}$$

The contraction of the primal (semilinear) problem is established in [25] and [22] based on Lemma 4.4, Lemma 4.5 and Theorem 4.7 as discussed above.

Theorem 4.8 (Contraction of the primal problem)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let u the solution to (1.2). Let $\theta \in (0,1]$, and let $\{\mathcal{T}_j, \mathbb{V}_j, u_j\}_{j\ge 0}$ be the sequence of meshes, finite element spaces and discrete solutions produced by GOAFEM. Then there exist constants $\gamma_p > 0$ and $0 < \alpha < 1$, depending on the initial mesh $\mathcal{T}_0$ and marking parameter θ such that*

$$|||u - u_{j+1}|||^2 + \gamma_p\eta_{j+1}^2 \le \alpha^2\left(|||u - u_j|||^2 + \gamma_p\eta_j^2\right). \tag{4.39}$$

5. CONTRACTION AND CONVERGENCE OF GOAFEM

In this section, we discuss the contraction and convergence of the GOAFEM described in §3. In particular, we show that the GOAFEM algorithm generates a sequence $\{\mathcal{T}_j, \mathbb{V}_j, u_j, \hat{z}_j\}_{j\ge 0}$ which contracts not only in the primal error as shown in §4, but also in a linear combination of the primal and limiting dual error. We emphasize that it would be difficult to derive convergence results in terms of problem (2.3) or (2.5), because at each refinement the problem is changing. So we show contraction in terms of the error in the limiting dual problem (2.6) as the target equation is fixed over the entire adaptive algorithm. Our approach to contraction in this section again relies on three main components: quasi-orthogonality, error-estimator as upper bound on error and estimator reduction. Here we discuss the relevant results for the limiting dual problem with an emphasis on those that differ significantly from the corresponding results for the primal problem. Note the limiting dual problem is not computable. We connect the error for the limiting dual problem to the computable quantities in the GOAFEM algorithm. For this purpose, we introduce Lemma 5.7, converting between limiting and approximate estimators in order to apply the Dörfler property to a computable quantity; and Lemma 5.8, bounding the discrete error between approximate and limiting dual solutions in terms of the primal error. We put these results together in Theorem 5.9 to establish the contraction of the combined quasi-error. Finally, the contraction of this form of the error is related to the error in the quantity of interest in Corollary 5.10.

5.1. **Quasi-orthogonality for Limiting-dual Problem.**

Similar to the proof of the quasi-orthogonality for the primal problem, we make use of an L_2-lifting argument for the limiting-dual problem. Let $\hat{z} \in H_0^1(\Omega)$ and $\hat{z}_1 \in \mathbb{V}_1$ be the solutions to (2.6) and (2.11), respectively. We again use the duality argument, and introduce the problem: Find $y \in H_0^1(\Omega)$ such that

$$a(y, v) + \langle b'(u)y, v\rangle = (\hat{z} - \hat{z}_1, v) \text{ for all } v \in H_0^1(\Omega) \tag{5.1}$$

Then we have the following L_2-lifting result for the limiting-dual problem.

Lemma 5.1 (Limiting-dual L_2-lifting)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let $\mathcal{T}_1$ be a conforming triangulation, and $\hat{z} \in H_0^1(\Omega)$ and $\hat{z}_1 \in \mathbb{V}_1$ be the solutions to (2.6) and (2.11), respectively. Assume that the solution y to (5.1) belongs to $H^{1+s}(\Omega) \cap H_0^1(\Omega)$ for some $0 < s \le 1$ such that*

$$|y|_{H^{1+s}(\Omega)} \le \bar{K}_R \|\hat{z} - \hat{z}_1\|_{L_2(\Omega)}. \tag{5.2}$$

Then

$$\|\hat{z} - \hat{z}_1\| \le \hat{C}_* h_0^s |\!|\!| \hat{z} - \hat{z}_1 |\!|\!|. \tag{5.3}$$

Proof. The proof follows that of Lemma 4.1. As in (4.12), we obtain for the limiting dual estimate (5.3)

$$\|\hat{z} - \hat{z}_1\|_{L_2} \le m_{\mathcal{E}}^{-1} \bar{K}_R \left(M_{\mathcal{E}} \bar{C}_{\mathcal{I}} + B \breve{C}_{\mathcal{I}} h_0 \right) h_0^s |\!|\!| \hat{z} - \hat{z}_1 |\!|\!|. \tag{5.4}$$

□

Remark 5.2. *To apply L_2-lifting to the difference between two Galerkin solutions, $\hat{z}_2 \in \mathbb{V}_2$ and $\hat{z}_1 \in V_1$ where $\mathcal{T}_2 \ge \mathcal{T}_1$, we use the same proof with (5.1) replaced by the problem: Find $y \in \mathbb{V}_2$ such that*

$$a(y, v) + \langle b'(u)y, v\rangle = \langle \hat{z}_2 - \hat{z}_1, v\rangle \textit{ for all } v \in \mathbb{V}_2. \tag{5.5}$$

With the help of Lemma 5.1, we obtain the quasi-orthogonality for the limiting-dual problem.

Lemma 5.3 (Quasi-orthogonality for Limiting Dual Problem)**.** *Let the problem data satisfy Assumption 2.1, and $\mathcal{T}_1, \mathcal{T}_2$ be two conforming triangulations with $\mathcal{T}_2 \ge \mathcal{T}_1$. Let $\hat{z} \in H_0^1(\Omega)$ the solution to (2.6) and $\hat{z}_i \in \mathbb{V}_i$ the solution to (2.11), $i = 1, 2$. There exists a constant $\hat{C}_* > 0$ depending on the problem data $\mathbf{D}$ and initial mesh $\mathcal{T}_0$, and a number $0 < s \le 1$ related to the regularity of (5.1), such that for sufficiently small h_0 we have*

$$|\!|\!| \hat{z} - \hat{z}_2 |\!|\!|^2 \le \hat{\Lambda} |\!|\!| \hat{z} - \bar{v} |\!|\!|^2 - |\!|\!| \hat{z}_2 - \bar{v} |\!|\!|^2, \quad \forall \bar{v} \in \mathbb{V}_2, \tag{5.6}$$

and in particular for $\bar{v} = \hat{z}_1$

$$|\!|\!| \hat{z} - \hat{z}_2 |\!|\!|^2 \le \hat{\Lambda}_G |\!|\!| \hat{z} - \hat{z}_1 |\!|\!|^2 - |\!|\!| \hat{z}_2 - \hat{z}_1 |\!|\!|^2 \tag{5.7}$$

where

$$\hat{\Lambda} := (1 - Bm_{\mathcal{E}}^{-1}\hat{C}_* h_0^s)^{-1} \text{ and } \hat{\Lambda}_G := (1 - B\hat{C}_*^2 h_0^{2s})^{-1}$$

and $\hat{C}_*$ is the constant from Lemma 5.1.

Proof. The proof follows same arguments as in Lemma 4.4, except that in place of the inequality in (4.18) we have for the limiting dual problem

$$a(u - u_2, v) + \langle b'(u)(\hat{z} - \hat{z}_2), v\rangle = 0 \text{ for all } v \in \mathbb{V}_2, \tag{5.8}$$

yielding

$$2a(\hat{z} - \hat{z}_2, \bar{v} - \hat{z}_2) \le 2B\|\hat{z} - \hat{z}_2\|_{L_2}\|\bar{v} - \hat{z}_2\|_{L_2}, \tag{5.9}$$

as in (4.19). The rest of the proof is similar to Lemma 4.4, and we omit it here. □

5.2. **Estimator Perturbations for Dual Sequence.** As we have seen in Theorem 4.7, the local Lipschitz property (cf. Lemma 4.6) plays a key role in deriving the estimator reduction property used to convert between estimators on different refinement levels in both the primal and limiting dual problems. The following lemma gives similar local Lipschitz properties for the approximate and limiting dual problems on a given refinement level.

Lemma 5.4 (Local Lipschitz Property for Dual Estimators)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let $\mathcal{T}$ be a conforming refinement of $\mathcal{T}_0$. Then for all $T \in \mathcal{T}$ and for any $v, w \in \mathbb{V}_{\mathcal{T}}$, it holds that*

$$|\zeta_{\mathcal{T},j}(v,T) - \zeta_{\mathcal{T},j}(w,T)| \le \bar{\Lambda}_1 \eta_{\mathcal{T}}(\mathbf{D}, T)\|v - w\|_{H^1(\omega_T)}. \tag{5.10}$$

In particular, for the error indicator of the limiting dual problem we have

$$|\zeta_{\mathcal{T}}(v,T) - \zeta_{\mathcal{T}}(w,T)| \le \bar{\Lambda}_1 \eta_{\mathcal{T}}(\mathbf{D}, T)\|v - w\|_{H^1(\omega_T)}. \tag{5.11}$$

The constant $\bar{\Lambda}_1 > 0$ depends on the dimension d and the regularity of the initial mesh $\mathcal{T}_0$.

Proof. The proof follows those in [8], [23] and is nearly identical to Lemma 4.6. It is sketched here. To prove (5.10), by (3.3) we have

$$\zeta_{\mathcal{T},j}^2(v,T) := h_T^2\|\hat{R}_j^*(v)\|_{L_2(T)}^2 + h_T\|J_T(v)\|_{L_2(\partial T)}^2, \quad v \in \mathbb{V}_{\mathcal{T}}. \tag{5.12}$$

Setting $e = v - w$ and applying linearity to the definition of the dual residual as given by (3.1), we obtain

$$\hat{R}_j^*(v) = g + \hat{\mathcal{L}}_j^*(w + e) = \hat{R}_j^*(w) + \hat{\mathcal{L}}_j^*(e).$$

By the same reasoning as (4.31), we get

$$\zeta_{\mathcal{T},j}(v,T) \le \zeta_{\mathcal{T},j}(w,T) + h_T\|\hat{\mathcal{L}}_j^*(e)\|_{L_2(T)} + h_T^{1/2}\|J(e)\|_{L_2(\partial T)}. \tag{5.13}$$

The term $\hat{\mathcal{L}}_j^*$ (respectively $\hat{\mathcal{L}}^*$ for the limiting dual) in (5.13) satisfies the same bound as the analogous term $\mathcal{D}$ in (4.31) of Lemma 4.6. Hence the bounds (5.10) and (5.11) hold with the same constants as in (4.29). □

With the help of Lemma 5.4, we are able to derive the following corollary, which addresses the error induced by switching between error indicators corresponding to the approximate and limiting dual problems on a given element.

Corollary 5.5. *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let $\mathcal{T}$ be a conforming refinement of $\mathcal{T}_0$, and u, u_j are the solutions to (1.2) and (2.9) problems, respectively. Let Θ and K_Z the constants given in Proposition 2.4. For all $T \in \mathcal{T}$ and for $v, w \in \mathbb{V}_\mathcal{T} \cap [z_-, z_+]$ the dual indicator on $\mathcal{T}$ satisfies*

$$|\zeta_{\mathcal{T},j}(v,T) - \zeta_{\mathcal{T},k}(w,T)| \le \bar{\Lambda}_1 \eta_\mathcal{T}(\mathbf{D},T)\|v-w\|_{H^1(\omega_T)} + \Theta K_Z h_T \|u_j - u_k\|_{L_2(T)}. \tag{5.14}$$

In particular, for $\mathcal{T} = \mathcal{T}_1$, we have for the limiting estimator

$$|\zeta_{1,1}(v,T) - \zeta_1(w,T)| \le \bar{\Lambda}_1 \eta_1(\mathbf{D},T)\|v-w\|_{H^1(\omega_T)} + \Theta K_Z h_T \|u - u_1\|_{L_2(T)}, \tag{5.15}$$

$$|\zeta_1(w,T) - \zeta_{1,1}(v,T)| \le \bar{\Lambda}_1 \eta_1(\mathbf{D},T)\|v-w\|_{H^1(\omega_T)} + \Theta K_Z h_T \|u - u_1\|_{L_2(T)}. \tag{5.16}$$

Proof. By the definition of the residuals for the approximate dual problems, for any $w \in \mathbb{V}_\mathcal{T}$ we have

$$\begin{aligned}\hat{R}^*_j(w) &= g + \nabla \cdot (A\nabla w) + b'(u_k)w + (b'(u_j) - b'(u_k))\, w \\ &= \hat{R}^*_k(w) + (b'(u_j) - b'(u_k))\, w.\end{aligned} \tag{5.17}$$

Using (5.17) in the definition of the dual indicator (3.3) and applying a generalized triangle inequality

$$\begin{aligned}\zeta_{\mathcal{T},j}(w,T) &= \left(h_T^2\|\hat{R}^*_k(w) + (b'(u_j) - b'(u_k))w\|^2_{L_2(T)} + h_T\|J_T(w)\|^2_{L_2(\partial T)}\right)^{1/2} \\ &\le \left(h_T^2\|\hat{R}^*_k(w)\|^2_{L_2(T)} + h_T\|J_T(w)\|^2_{L_2(\partial T)}\right)^{1/2} + h_T\|b'(u_j) - b'(u_k)w\|_{L_2(T)} \\ &\le \zeta_{\mathcal{T},k}(w,T) + \Theta K_Z h_T \|u_j - u_k\|_{L_2(T)}.\end{aligned} \tag{5.18}$$

Applying (5.10) in Lemma 5.4 to the estimate (5.18), we obtain (5.14). ☐

As an immediate consequence of Corollary 5.5, we have the following results on the error induced by switching between dual estimators over a collection of elements on a given refinement level. This estimate plays a key role in the contraction argument below, as we apply it to switching between the estimator for the limiting dual and the computed error estimators for the approximate dual problems in the GOAFEM algorithm.

Corollary 5.6. *Let the hypotheses of Corollary 5.5 hold. Then for any subsets $\mathcal{M}_1, \mathcal{M}_2 \subseteq \mathcal{T}_1$ and arbitrary $\delta_1, \delta_2, \delta_A, \delta_B > 0$*

$$\begin{aligned}\zeta_1^2(v,\mathcal{M}_1) \ge{}& (1+\delta_1)^{-1}(1+\delta_A)^{-1}\zeta_{1,1}^2(w,\mathcal{M}_1) \\ &- (1+\delta_1)^{-1}\delta_A^{-1}\Theta^2 K_Z^2 h_0^2\|u-u_1\|^2_{L_2} - (d+2)\delta_1^{-1}\bar{\Lambda}_1^2\eta_0^2\|v-w\|^2_{H^1}\end{aligned} \tag{5.19}$$

$$\begin{aligned}\zeta_{1,1}^2(w,\mathcal{M}_2) \ge{}& (1+\delta_2)^{-1}(1+\delta_B)^{-1}\zeta_1^2(v,\mathcal{M}_2) \\ &- (1+\delta_2)^{-1}\delta_B^{-1}\Theta^2 K_Z^2 h_0^2\|u-u_1\|^2_{L_2} - (d+2)\delta_2^{-1}\bar{\Lambda}_1^2\eta_0^2\|v-w\|^2_{H^1}.\end{aligned} \tag{5.20}$$

Proof. The conclusions follow by squaring inequality (5.15) (respectively (5.16)), applying Young's inequality twice, and then summing over element $T \in \mathcal{M}_1$ (respectively $T \in \mathcal{M}_2$). The H^1 norm is summed over all elements $T \in \mathcal{T}_1$ counting each element $d+2$ times, the maximum number of elements in each patch ω_T. ☐

5.3. **Contraction of GOAFEM.**

The main contraction argument Theorem 5.9 follows after two more lemmas. The first combines a sequence of estimates to convert the non-computable limiting estimator for the dual problem to a computable quantity, apply the Dörfler property and then convert back. The second relates the difference between the Galerkin solutions of the limiting and approximate dual problems to the primal error. Motivated by estimator reduction for the limiting dual problem as in equation (4.38)

$$\zeta_2^2(\hat{z}_2, \mathcal{T}_2) \leq (1+\delta)\left\{\zeta_1^2(\hat{z}_1, \mathcal{T}_1) - \lambda\zeta_1^2(\hat{z}_1, \mathcal{M})\right\} + (1+\delta^{-1})\Lambda_1\eta_0^2 |||\hat{z}_2 - \hat{z}_1|||^2 \tag{5.21}$$

the following lemma addresses the conversion between the limiting estimator $\zeta_1^2(\hat{z}_1, \mathcal{M})$ and and the computable estimator $\zeta_{1,1}^2(\hat{z}_1^1, \mathcal{M})$ necessary for marking the mesh for refinement.

Lemma 5.7. *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let Θ and K_Z as given by Proposition 2.4, C_* as given by Lemma 4.1 and Λ_1 as given in Lemma 4.7. Let*

u the solution to (1.2), u_1 the solution to (2.9),
$\hat{z}$ the solution to (2.6), $\hat{z}_1$ the solution to (2.11) $\hat{z}_1^1$ the solution to (2.10).

Let $\zeta_{1,1}(\hat{z}_1^1, \mathcal{M})$ satisfy the Dörfler property for $\mathcal{M} \subset \mathcal{T}_1$, namely $\zeta_{1,1}^2(\hat{z}_1^1, \mathcal{M}) \geq \theta^2\zeta_{1,1}^2(\hat{z}_1^1, \mathcal{T}_1)$. Then for arbitrary $\delta_1, \delta_2, \delta_A, \delta_B > 0$ there is a $\delta_4 > 0$ such that

$$\begin{aligned} -\zeta_1^2(\hat{z}_1, \mathcal{M}) \leq & -\frac{\beta\theta^2}{(1+\delta_4)}\zeta_1^2(\hat{z}_1, \mathcal{T}_1) - \frac{(1-\beta)\theta^2}{(1+\delta_4)C_1^2}|||\hat{z} - \hat{z}_1|||^2 \\ & + \left(\frac{\theta^2}{(1+\delta_A)(1+\delta_2)\delta_B} + \frac{1}{\delta_A}\right)\frac{\Theta^2 K_Z^2 C_*^2 h_0^{2(1+s)}}{(1+\delta_1)}|||u - u_1|||^2 \\ & + \left(\frac{\theta^2}{(1+\delta_1)(1+\delta_A)\delta_2} + \frac{1}{\delta_1}\right)\Lambda_1\eta_0^2(\mathbf{D}, \mathcal{T}_0)|||\hat{z}_1 - \hat{z}_1^1|||^2. \end{aligned} \tag{5.22}$$

Proof. From Corollary 5.6, L_2-lifting 4.1 and coercivity (2.2)

$$\begin{aligned} -\zeta_1^2(\hat{z}_1, \mathcal{M}) \leq & -(1+\delta_1)^{-1}(1+\delta_A)^{-1}\zeta_{1,1}^2(\hat{z}_1^1, \mathcal{M}) \\ & + (1+\delta_1)^{-1}\delta_A^{-1}\Theta^2 K_Z^2 h_0^2\|u - u_1\|_{L_2}^2 + \delta_1^{-1}\bar{\Lambda}_1^2(d+2)\eta_0^2\|\hat{z}_1 - \hat{z}_1^1\|_{H^1}^2 \\ \leq & -(1+\delta_1)^{-1}(1+\delta_A)^{-1}\zeta_{1,1}^2(\hat{z}_1^1, \mathcal{M}) \\ & + (1+\delta_1)^{-1}\delta_A^{-1}\Theta^2 K_Z^2 C_*^2 h_0^{2(1+s)}|||u - u_1|||^2 + \delta_1^{-1}\Lambda_1\eta_0^2|||\hat{z}_1 - \hat{z}_1^1|||^2 \end{aligned} \tag{5.23}$$

with $\Lambda_1 := \bar{\Lambda}_1^2(d+2)m_{\mathcal{E}}^{-2}$. The Dörfler property may be applied to the first term on the RHS of (5.23)

$$-\zeta_{1,1}^2(\hat{z}_1^1, \mathcal{M}) \leq -\theta^2\zeta_{1,1}^2(\hat{z}_1^1). \tag{5.24}$$

Converting back to he limiting estimator by (5.20) in Corollary 5.6

$$\begin{aligned} -\zeta_{1,1}^2(\hat{z}_1^1) \leq & -(1+\delta_2)^{-1}(1+\delta_B)^{-1}\zeta_1^2(\hat{z}_1, \mathcal{M}) \\ & + (1+\delta_2)^{-1}\delta_B^{-1}\Theta^2 K_Z^2 C_*^2 h_0^{2(1+s)}|||u - u_1|||^2 + \delta_2^{-1}\Lambda_1\eta_0^2|||\hat{z}_1 - \hat{z}_1^1|||^2. \end{aligned} \tag{5.25}$$

Define δ_4 by

$$(1+\delta_4) := (1+\delta_1)(1+\delta_2)(1+\delta_A)(1+\delta_B). \tag{5.26}$$

Then by plugging (5.24) and (5.25) in the first term on the RHS of (5.23), we obtain

$$\begin{aligned}-\zeta_1^2(\hat{z}_1,\mathcal{M}) &\le -\theta^2(1+\delta_4)^{-1}\zeta_1^2(\hat{z}_1)\\ &\quad + \left(\theta^2(1+\delta_A)^{-1}(1+\delta_2)^{-1}\delta_B^{-1}+\delta_A^{-1}\right)(1+\delta_1)^{-1}\Theta^2K_Z^2C_*^2h_0^{2(1+s)}|\!|\!|u-u_1|\!|\!|^2\\ &\quad + \left(\theta^2(1+\delta_1)^{-1}(1+\delta_A)^{-1}\delta_2^{-1}+\delta_1^{-1}\right)\Lambda_1\eta_0^2|\!|\!|\hat{z}_1-\hat{z}_1^1|\!|\!|^2. \qquad (5.27)\end{aligned}$$

Finally, we split the first term on the RHS of (5.27) into two pieces for some $\beta \in (0,1)$, and apply the upper-bound estimate (4.25) in Lemma 4.5 to the second piece yielding

$$\begin{aligned}-\zeta_1^2(\hat{z}_1,\mathcal{M}) &\le -\beta\theta^2(1+\delta_4)^{-1}\zeta_1^2(\hat{z}_1) - (1-\beta)\theta^2(1+\delta_4)^{-1}C_1^{-2}|\!|\!|\hat{z}-\hat{z}_1|\!|\!|^2\\ &\quad + \left(\theta^2(1+\delta_A)^{-1}(1+\delta_2)^{-1}\delta_B^{-1}+\delta_A^{-1}\right)(1+\delta_1)^{-1}\Theta^2K_Z^2C_*^2h_0^{2(1+s)}|\!|\!|u-u_1|\!|\!|^2\\ &\quad + \left(\theta^2(1+\delta_1)^{-1}(1+\delta_A)^{-1}\delta_2^{-1}+\delta_1^{-1}\right)\Lambda_1\eta_0^2|\!|\!|\hat{z}_1-\hat{z}_1^1|\!|\!|^2.\end{aligned}$$

This completes the proof. ☐

We may convert $|\!|\!|\hat{z}_1-\hat{z}_1^1|\!|\!|$ in the last term on the RHS of (5.22) to the error $|\!|\!|u-u_1|\!|\!|$ as stated in the following lemma.

Lemma 5.8. *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let Θ and K_Z the constants given in Proposition 2.4 and C_* and $\hat{C}_*$ the constants given by Lemmas 4.1 and 5.1, respectively. Let*

u *the solution to* (1.2), u_1 *the solution to* (2.9),

$\hat{z}_1$ *the solution to* (2.11), $\hat{z}_1^1$ *the solution to* (2.10).

Then

$$|\!|\!|\hat{z}_1-\hat{z}_1^1|\!|\!| \le \Theta K_Z C_*\hat{C}_* h_0^{2s}|\!|\!|u-u_1|\!|\!|. \qquad (5.28)$$

Proof. Recall that

$$\begin{aligned}&\hat{z}_1 \text{ solves } a(\hat{z}_1,v)+\langle b'(u)\hat{z}_1,v\rangle = g(v), \quad \text{for all } v\in\mathbb{V}_1, &(5.29)\\ &\hat{z}_1^1 \text{ solves } a(\hat{z}_1^1,v)+\langle b'(u_1)\hat{z}_1^1,v\rangle = g(v), \quad \text{for all } v\in\mathbb{V}_1. &(5.30)\end{aligned}$$

Subtracting (5.30) from (5.29) and rearranging terms, we get

$$a(\hat{z}_1-\hat{z}_1^1,v)+\langle (b'(u)-b'(u_1))\hat{z}_1,v\rangle = \langle b'(u_1)(\hat{z}_1^1-\hat{z}_1),v\rangle,\ v\in\mathbb{V}_1. \qquad (5.31)$$

In particular, for $v=\hat{z}_1-\hat{z}_1^1\in\mathbb{V}_1$ equation (5.31) yields

$$\begin{aligned}|\!|\!|\hat{z}_1-\hat{z}_1^1|\!|\!|^2 &= -\langle (b'(u)-b'(u_1))\hat{z}_1,\hat{z}_1-\hat{z}_1^1\rangle - \langle b'(u_1)(\hat{z}_1-\hat{z}_1^1),\hat{z}_1-\hat{z}_1^1\rangle\\ &\le -\langle (b'(u)-b'(u_1))\hat{z}_1,\hat{z}_1-\hat{z}_1^1\rangle \qquad (5.32)\end{aligned}$$

where in the last inequality, we used the monotonicity assumption of b in Assumption (2.1). Now applying the Lipschitz property of b', the *a priori* L_∞ bounds on the dual solution $\hat{z}_1$ (cf. Proposition 2.4), and both primal and dual L_2 lifting in (5.32), we obtain

$$\begin{aligned}|\!|\!|\hat{z}_1-\hat{z}_1^1|\!|\!|^2 &\le \Theta K_Z\|u-u_1\|_{L_2}\|\hat{z}_1-\hat{z}_1^1\|_{L_2}\\ &\le \Theta K_Z C_*\hat{C}_* h_0^{2s}|\!|\!|u-u_1|\!|\!|\,|\!|\!|\hat{z}_1-\hat{z}_1^1|\!|\!| \qquad (5.33)\end{aligned}$$

from which the result follows. ☐

Now we are in position to show the contraction of GOAFEM in terms of the combined quasi-error which is a linear combination of the energy errors and error estimators in primal and limiting dual problems.

Theorem 5.9 (Contraction of GOAFEM)**.** *Let the problem data satisfy Assumption 2.1 and Assumption 2.2. Let*

u *the solution to* (1.2), u_j *the solution to* (2.9),
$\hat{z}$ *the solution to* (2.6), $\hat{z}_j$ *the solution to* (2.11).

Let $\theta \in (0,1]$, *and let* $\{\mathcal{T}_j, \mathbb{V}_j\}_{j\geq 0}$ *be the sequence of meshes and finite element spaces produced by GOAFEM. Let* $\gamma_p > 0$ *as given by Theorem 4.8. Then for sufficient small mesh size* h_0, *there exist constants* $\gamma > 0, \pi > 0$ *and* $\alpha_D \in (0,1)$ *such that*

$$\begin{aligned} &|\!|\!| \hat{z} - \hat{z}_2 |\!|\!|^2 + \gamma \zeta_2^2(\hat{z}_2) + \pi |\!|\!| u - u_2 |\!|\!|^2 + \pi\gamma_p \eta_2^2(u_2) \\ &\leq \alpha_D^2 \left(|\!|\!| \hat{z} - \hat{z}_1 |\!|\!|^2 + \gamma \zeta_1^2(\hat{z}_1) + \pi |\!|\!| u - u_1 |\!|\!|^2 + \pi\gamma_p \eta_1^2(u_1) \right). \end{aligned} \tag{5.34}$$

Proof. For simplicity, we denote $\eta_0 = \eta_0(\mathbf{D}, \mathcal{T}_0)$ and $\zeta_k(\hat{z}_k) = \zeta_k(\hat{z}_k, \mathcal{T}_k),\ k = 1, 2$. By the estimator reduction for the limiting dual problem (4.38), for arbitrary $\delta > 0$ we have

$$\zeta_2^2(\hat{z}_2) \leq (1+\delta)\left\{ \zeta_1^2(\hat{z}_1) - \lambda \zeta_1^2(\hat{z}_1, \mathcal{M}) \right\} + (1+\delta^{-1})\Lambda_1 \eta_0^2 |\!|\!| \hat{z}_2 - \hat{z}_1 |\!|\!|^2, \tag{5.35}$$

where $\lambda = 1 - 2^{-1/d}$. Recall the quasi-orthogonality estimate in the limiting dual problem from Lemma 5.3

$$|\!|\!| \hat{z} - \hat{z}_2 |\!|\!|^2 \leq \hat{\Lambda}_G |\!|\!| \hat{z} - \hat{z}_1 |\!|\!|^2 - |\!|\!| \hat{z}_2 - \hat{z}_1 |\!|\!|^2. \tag{5.36}$$

Adding (5.36) to a positive multiple γ (to be determined) of (5.35) and applying the results of Lemmas 5.7 and 5.8 obtain

$$\begin{aligned} |\!|\!| \hat{z} - \hat{z}_2 |\!|\!|^2 + \gamma \zeta_2^2(\hat{z}_2) \leq A |\!|\!| \hat{z} - \hat{z}_1 |\!|\!|^2 + \gamma M \zeta_1^2(\hat{z}_1) + D |\!|\!| u - u_1 |\!|\!|^2 \\ + \left(\gamma (1+\delta^{-1}) \Lambda_1 \eta_0^2 - 1 \right) |\!|\!| \hat{z}_2 - \hat{z}_1 |\!|\!|^2. \end{aligned} \tag{5.37}$$

We first set $\gamma := (1+\delta^{-1})^{-1} \Lambda_1^{-1} \eta_0^{-2}$ to eliminate the last term in (5.37). This yields

$$|\!|\!| \hat{z} - \hat{z}_2 |\!|\!|^2 + \gamma \zeta_2^2(\hat{z}_2) \leq A |\!|\!| \hat{z} - \hat{z}_1 |\!|\!|^2 + \gamma M \zeta_1^2(\hat{z}_1) + D |\!|\!| u - u_1 |\!|\!|^2, \tag{5.38}$$

where the coefficients A and M of (5.38) are given by

$$A = \hat{\Lambda}_G - (1-\beta)\lambda\theta^2 \delta (1+\delta_4)^{-1} C_1^{-2} \Lambda_1^{-1} \eta_0^{-2} \tag{5.39}$$

$$M = (1+\delta)(1 - \beta\lambda\theta^2 (1+\delta_4)^{-1}) \tag{5.40}$$

where δ_4 satisfies $(1+\delta_4) := (1+\delta_1)(1+\delta_2)(1+\delta_A)(1+\delta_B)$ as was given in (5.26).

For contraction, we require $A < 1$ and $M < 1$ for the coefficients defined by (5.39) and (5.40), that is, we need to choose a $\beta \in (0,1)$ such that

$$\frac{\delta}{1+\delta} \frac{1+\delta_4}{\lambda\theta^2} < \beta < 1 - \frac{(\hat{\Lambda}_G - 1)\Lambda_C}{\delta} \frac{1+\delta_4}{\lambda\theta^2}, \tag{5.41}$$

with $\Lambda_C := C_1^2 \Lambda_1 \eta_0^2$. To demonstrate the existence of such a β, set

$$\delta_4 = \delta = \frac{1}{2}\lambda\theta^2. \tag{5.42}$$

Then we require the mesh size h_0 sufficiently small, such that

$$\hat{\Lambda}_G < 1 + \frac{\lambda^2\theta^4}{2(2+\lambda\theta^2)\Lambda_C}, \tag{5.43}$$

for the given $\theta \in (0,1)$. Note the conditions (5.42) and (5.43) guarantee that the interval in (5.41) is nonempty, so there exists a β such that

$$\frac{1}{2} < \beta < 1 - \frac{(\hat{\Lambda}_G - 1)\Lambda_C}{\lambda\theta^2} \left(1 + \frac{2}{\lambda\theta^2} \right).$$

It remains to control the last term in (5.38). For simplicity, we assume $\delta_1 = \delta_2 = \delta_A = \delta_B =: \delta_C$. Then the coefficient D in (5.38) is given by

$$D = \delta\lambda\Theta^2 K_Z^2 C_*^2 h_0^{2s}\left(\frac{\theta^2 + (1+\delta_C)^2}{(1+\delta_C)^2\delta_C}\right)\left(\frac{h_0^2}{\Lambda_1\eta_0^2(1+\delta_C)} + \hat{C}_*^2 h_0^{2s}\right). \tag{5.44}$$

To control the primal error term with the coefficient D as given by (5.44), we add a positive multiple π (to be determined) of the primal contraction result (4.39) of Theorem 4.8 to (5.39) yieding

$$\begin{aligned}&|\!|\!|\hat{z} - \hat{z}_2|\!|\!|^2 + \gamma\zeta_2^2(\hat{z}_2) + \pi|\!|\!|u - u_2|\!|\!|^2 + \pi\gamma_p\eta_2^2(u_2)\\ &\le A|\!|\!|\hat{z} - \hat{z}_1|\!|\!|^2 + \gamma M\zeta_1^2(\hat{z}_1) + (D + \alpha^2\pi)|\!|\!|u - u_1|\!|\!|^2 + \alpha^2\pi\gamma_P\eta_1^2(u_1).\end{aligned} \tag{5.45}$$

We choose π to ensure $D + \alpha^2\pi < \pi$, namely,

$$\pi > \frac{D}{1-\alpha^2} \tag{5.46}$$

and set

$$\alpha_D^2 := \max\left\{A, M, \frac{D+\alpha^2\pi}{\pi}, \alpha^2\right\} < 1. \tag{5.47}$$

Then the combined quasi-error satisfies the contraction property (5.34). □

For simplicity, we denote by

$$\bar{Q}^2(u_j, \hat{z}_j) = |\!|\!|\hat{z} - \hat{z}_j|\!|\!|^2 + \gamma\zeta_j^2(\hat{z}_j) + \pi|\!|\!|u - u_j|\!|\!|^2 + \pi\gamma_p\eta_j^2(u_j)$$

the combined quasi-error in (5.34). The following corollary gives the contraction of the error in the goal function, which is determined by the contraction of the combined quasi-error.

Corollary 5.10. *Let the assumptions in Theorem 5.9 hold. Then the error in the goal function is controlled by a constant multiple of the square of the combined quasi-error, i.e.,*

$$|g(u) - g(u_j)| \le C\bar{Q}_j^2(u_j, \hat{z}_j) \le \alpha_D^{2j} C\bar{Q}_0^2(u_0, \hat{z}_0). \tag{5.48}$$

Proof. Choosing the test function $v = u - u_j$ in (2.6), and by linearity and Galerkin orthogonality for the primal problem, we obtain

$$\begin{aligned}g(u) - g(u_j) &= a(\hat{z}, u) + \langle b'(u)\hat{z}, u\rangle - a(\hat{z}, u_j) - \langle b'(u)\hat{z}, u_j\rangle\\ &= a(u - u_j, \hat{z}) + \langle b'(u)(u - u_j), \hat{z}\rangle\\ &= a(u - u_j, \hat{z}) + \langle \mathcal{B}_j(u - u_j), \hat{z}\rangle + \langle (b'(u) - \mathcal{B}_j)(u - u_j), \hat{z}\rangle\\ &= a(u - u_j, \hat{z} - \hat{z}_j) + \langle b(u) - b(u_j), \hat{z} - \hat{z}_j\rangle + \langle (b'(u) - \mathcal{B}_j)(u - u_j), \hat{z}\rangle.\end{aligned} \tag{5.49}$$

The third term in the last line of (5.49) represents the error induced by switching from (2.6) to (2.3). This term may be bounded in terms of the constants and L_∞ estimates in Proposition 2.4 and

$$\|b'(u) - \mathcal{B}_j\| = \left\|\int_0^1 b'(u) - b'\left(u_j + \xi(u - u_j)\right) d\xi\right\| \le \frac{\Theta}{2}\|u - u_j\|,$$

yielding

$$\begin{aligned}\langle (b'(u) - \mathcal{B}_j)(u - u_j), \hat{z}\rangle &\le K_Z\|b'(u) - \mathcal{B}_j\|_{L_2}\|u - u_j\|_{L_2}\\ &\le \frac{1}{2}\Theta K_Z\|u - u_j\|_{L_2}^2.\end{aligned} \tag{5.50}$$

Then by (5.49), (5.50), the Cauchy-Schwarz inequality and L_2-lifting as in Lemmas 4.1 and 5.1

$$\begin{aligned}|g(u)-g(u_j)| &\le |||u-u_j|||\,|||\hat z-\hat z_j||| + B\|u-u_j\|_{L_2}\|\hat z-\hat z_j\|_{L_2} + \frac{1}{2}\Theta K_Z\|u-u_j\|_{L_2}^2\\ &\le (1+BC_*\hat C_* h_0^{2s})|||u-u_j|||\,|||\hat z-\hat z_j||| + \frac{1}{2}\Theta K_Z C_*^2 h_0^{2s}|||u-u_j|||^2\\ &\le \frac{1}{2}\left(1+(\Theta K_Z C_* + B\hat C_*)C_* h_0^{2s}\right)|||u-u_j|||^2 + \frac{1}{2}(1+BC_*\hat C_* h_0^{2s})|||\hat z-\hat z_j|||^2.\end{aligned} \tag{5.51}$$

Therefore the error in the goal function is bounded above by a constant multiple of the square of the combined quasi-error $\bar{Q}^2(u_j, \hat{z}_j)$. Thus (5.48) follows by the contraction result in Theorem 5.9. ☐

6. NUMERICAL EXPERIMENTS

In this section, we present some numerical experiments implemented using FETK [20], which is a fairly standard set of finite element modeling libraries for approximating the solutions to systems of nonlinear elliptic and parabolic equations. In these experiments, we consider $\Omega = [0,1]^2$ and try to solve the following model problem:

$$\mathcal{N}(u) := -\Delta u + 3u^3 = f, \tag{6.1}$$

with homogeneous Dirichlet boundary condition. Here the source function f is chosen such that the exact solution is given by

$$u(x,y) = \frac{\sin(\pi x)\sin(\pi y)}{2(x-0.5)^2 + 2(y-0.5)^2 + 10^{-3}}.$$

We solve the nonlinear problem using both the AFEM and the GOAFEM algorithms. Figure 1 shows a typical adaptive mesh as well as the solution of this semilinear equation.



FIGURE 1. The mesh and finite element solution to the model problem after 12 GOAFEM iterations.

In the adaptive algorithms, we use the Dörfler marking strategies: (3.7) for the primal problem and (3.8) for the dual problem, with the same parameter $\theta = 0.4$. For the primal nonlinear problem (6.1), at each refinement we use a Newton-type iteration to solve the resulting nonlinear system of algebraic equations, which reduces the nonlinear residual



FIGURE 2. The reduction rate in the goal error $|g(u-u_h)|$.

to the tolerance $\|F(u)\|_{L_2} \leq 10^{-7}$. On the initial triangulation, we use a zero initial guess for the Newton iteration; then for each subsequent refinement, we interpolate the numerical solution from the previous step to the current triangulation and then use it as the initial guess for the Newton iteration. By doing this, we have a good initial guess for the Newton iteration so that one could expect a quadratic convergence rate of the nonlinear iterations. In fact, according to our numerical experiments, it usually takes 4 or 5 Newton iterations to reach the setting tolerance.

To test the performance of the GOAFEM algorithm, we take the goal function

$$g = 100e^{-100((x-0.5)^2+(y-0.5)^2)}, \tag{6.2}$$

so that it captures the singularity of the solution to (6.1). Thus the error in goal functional $|g(u-u_h)|$ gives us an error by weighted average. Figure 2 shows the reduction rate of the goal error $|g(u-u_h)|$ for the GOAFEM algorithm. The oscillation at the first a few iterations in Figure 2 reflects the fact that the mesh size is not small enough, which is one of the requirements in our theory. But after a few iterations, the goal error reduces at certain rate, which was predicted by Theorem 5.48. This result confirms our theory.

For comparison with the standard AFEM algorithm, we also show in Figure 3 the error reduction in H^1 semi-norm $|u-u_h|_{H^1}$ for both AFEM and GOAFEM algorithms with the same goal function given in (6.2). As one can see from Figure 3, reducing H^1 error to the same magnitude with both algorithms, the GOAFEM algorithm takes about 5 fewer refinement steps than the standard AFEM algorithm.

7. CONCLUSION

In this article we developed convergence theory for a class of goal-oriented adaptive finite element algorithms for second order semilinear elliptic equations. We first introduced several approximate dual problems, and briefly discussed the target problem class. We then reviewed some standard facts concerning conforming finite element discretization and error-estimate-driven adaptive finite element methods (AFEM). We included a brief summary of *a priori* estimates for semilinear problems, and then described goal-oriented variations of the standard approach to AFEM (GOAFEM). Following the recent work of Mommer-Stevenson and Holst-Pollock for linear problems, we established contraction of GOAFEM for the primal problem. We also developed some additional



FIGURE 3. The reduction rate in the H^1 norm of the error $u - u_h$.

estimates that make it possible to establish contraction of the combined quasi-error, and showed convergence in the sense of the quantity of interest. Some simple numerical experiments confirmed these theoretical predictions. Our analysis was based on the recent contraction frameworks for the semilinear problem developed by Holst, Tsogtgerel, and Zhu and Bank, Holst, Szypowski and Zhu and those for linear problems as in Cascon, Kreuzer, Nochetto and Siebert, and Nochetto, Siebert, and Veeser. In addressing the goal-oriented problem we based our approach on that of Mommer and Stevenson for symmetric linear problems and Holst and Pollock for nonsymmetric problems. However, unlike the linear case, we were faced with tracking linearized and approximate dual sequences in order to establish contraction with respect to the quantity of interest.

In the present paper we assume the primal and approximate dual solutions are solved on the same mesh at each iteration. The determination of strong convergence results for a method which solves the primal (nonlinear) problem on a coarse mesh and the dual on a fine mesh is the subject of future investigation.

ACKNOWLEDGMENTS

MH was supported in part by NSF Awards 0715146 and 0915220, and by DOD/DTRA Award HDTRA-09-1-0036. SP was supported in part by NSF Award 0715146. YZ was supported in part by NSF Award 0715146 and by DOD/DTRA Award HDTRA-09-1-0036.

REFERENCES

[1] O. Axelsson and V. A. Barker. *Finite element solution of boundary value problems: theory and computation*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 2001.

[2] R. Bank, M. Holst, R. Szypowski, and Y. Zhu. Convergence of AFEM for semilinear problems with inexact solvers, 2011.

[3] R. Bank, M. Holst, R. Szypowski, and Y. Zhu. Finite element error estimates for critical growth semilinear problems without angle conditions, 2011.

[4] R. Becker and R. Rannacher. A feed-back approach to error control in finite element methods: Basic analysis and examples. *East-West Journal of Numerical Mathematics*, 4:237–264, 1996.

[5] R. Becker and R. Rannacher. An optimal control approach to a posteriori error estimation in finite element methods. *Acta Numerica*, pages 1–102, 2001.

[6] P. Binev, W. Dahmen, and R. DeVore. Adaptive finite element methods with convergence rates. *Numer. Math.*, 97(2):219–268, 2004.

[7] S. Brenner and L. Scott. *The Mathematical Theory of Finite Element Methods*. Springer-Verlag, third edition, 2008.

[8] J. M. Cascon, C. Kreuzer, R. H. Nochetto, and K. G. Siebert. Quasi-optimal convergence rate for an adaptive finite element method. *SIAM J. Numer. Anal.*, 46(5):2524–2550, 2008.

[9] P. G. Ciarlet. *Finite Element Method for Elliptic Problems*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 2002.

[10] W. Dahmen, A. Kunoth, and J. Vorloeper. *Convergence of Adaptive Wavelet Methods for Goal-oriented Error Estimation*. Sonderforschungsbereich 611, Singuläre Phänomene und Skalierung in Mathematischen Modellen. SFB 611, 2006.

[11] W. Dörfler. A convergent adaptive algorithm for Poisson's equation. *SIAM Journal on Numerical Analysis*, 33:1106–1124, 1996.

[12] K. Eriksson, D. Estep, P. Hansbo, and C. Johnson. Introduction to adaptive methods for differential equations. *Acta Numerica*, pages 105–158, 1995.

[13] D. Estep, M. Holst, and M. Larson. Generalized green's functions and the effective domain of influence. *SIAM J. Sci. Comput*, 26:1314–1339, 2002.

[14] D. Estep, M. Holst, and D. Mikulencak. Accounting for stability: A posteriori error estimates based on residuals and variational analysis. In *Communications in Numerical Methods in Engineering*, pages 200–2, 2001.

[15] D. Estep, M. G. Larson, and R. D. Williams. Estimating the error of numerical solutions of systems of reaction-diffusion equations. *Mem. Amer. Math. Soc.*, 146(696):101–109, 2000.

[16] L. C. Evans. *Partial Differential Equations (Graduate Studies in Mathematics, V. 19) GSM/19*. American Mathematical Society, 1998.

[17] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Springer-Verlag, 1977.

[18] M. Giles and E. Süli. Adjoint methods for PDEs: a posteriori error analysis and postprocessing by duality. *Acta Numerica*, 11:145–236, 2003.

[19] T. Grätsch and K.-J. Bathe. A posteriori error estimation techniques in practical finite element analysis. *Computers & Structures*, 83(4-5):235 – 265, 2005.

[20] M. Holst. Adaptive numerical treatment of elliptic systems on manifolds. 15(1–4):139–191, 2001. Available as arXiv:1001.1367 [math.NA].

[21] M. Holst. Applications of domain decomposition and partition of unity methods in physics and geometry, 2003.

[22] M. Holst, J. McCammon, Z. Yu, Y. Zhou, and Y. Zhu. Adaptive finite element modeling techniques for the Poisson-Boltzmann equation. *Communications in Computational Physics*, 11(1):179–214, 2012. Available as arXiv:1009.6034 [math.NA].

[23] M. Holst and S. Pollock. Convergence of goal oriented methods for nonsymmetric problems, 2011.

[24] M. Holst, R. Szypowski, and Y. Zhu. Two-grid methods for semilinear interface problems, 2012.

[25] M. Holst, G. Tsogtgerel, and Y. Zhu. Local and global convergence of adaptive methods for nonlinear partial differential equations, 2008.

[26] A. Jüngel and A. Unterreiter. Discrete minimum and maximum principles for finite element approximations of non-monotone elliptic equations. *Numer. Math.*, 99(3):485–508, 2005.

[27] J. Karatson and S. Korotov. Discrete maximum principles for finite element solutions of nonlinear elliptic problems with mixed boundary conditions. *Numerische Mathematik*, 99:669–698, 2005.

[28] T. Kerkhoven and J. W. Jerome. L_∞ stability of finite element approximations of elliptic gradient equations. *Numerische Mathematik*, 57:561–575, 1990.

[29] S. Kesavan. *Topics in Functional Analysis and Applications*. John Wiley and Sons, Inc., New York, NY, 1989.

[30] S. Korotov. A posteriori error estimation of goal-oriented quantities for elliptic type bvps. *Journal of Computational and Applied Mathematics*, 191(2):216 – 227, 2006.

[31] K. Mekchay and R. Nochetto. Convergence of adaptive finite element methods for general second order linear elliptic PDE. *SINUM*, 43(5):1803–1827, 2005.

[32] M. S. Mommer and R. Stevenson. A goal-oriented adaptive finite element method with convergence rates. *SIAM J. Numer. Anal.*, 47(2):861–886, 2009.

[33] K.-S. Moon, E. von Schwerin, A. Szepessy, and R. Tempone. Convergence rates for an adaptive dual weighted residual finite element algorithm. *BIT*, 46(2):367–407, 2006.

[34] R. H. Nochetto, K. G. Siebert, and A. Veeser. *Theory of adaptive finite element methods: an introduction*, pages 409 – 542. Springer, 2009.

[35] J. Oden and S. Prudhomme. Goal-oriented error estimation and adaptivity for the finite element method. *Computers and Mathematics with Applications*, 41:735–756, 2001.

[36] E. G. Sewell. Automatic generation of triangulations for piecewise polynomial approximation. In *Ph. D. dissertation*. Purdue Univ., West Lafayette, Ind., 1972.

[37] G. Strang and G. J. Fix. *An Analysis of the Finite Element Method*. Prentice-Hall (Series in Automatic Computation), Englewood Cliffs, N. J., 1973.

[38] M. Struwe. *Variational Methods*. Springer-Verlag, Berlin, Germany, 3 edition, 2000.

[39] R. Verfürth. A posteriori error estimates for nonlinear problems. finite element discretizations of elliptic equations. *Mathematics of Computation*, 62(206):445–475, Apr. 1994.

[40] R. Verfürth. *A review of a posteriori error estimation and adaptive mesh refinement tecniques*. B. G. Teubner, 1996.

E-mail address: `mholst@math.ucsd.edu`

E-mail address: `snpolloc@math.ucsd.edu`

E-mail address: `zhu@math.ucsd.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA SAN DIEGO, LA JOLLA CA 92093